

Case Report

Coexistence of Recidivans, Mucocutaneous and Osseous Leishmaniasis in an Immunocompromised Patient

Hira Tariq¹, Shahbaz Aman², Ambereen Anwar³, Ameena Ashraf⁴

^{1,2}Department of Dermatology, SIMS/ Services Hospital, Lahore; ^{3,4}Department of Pathology, Allama Iqbal Medical College, Lahore

Abstract

Leishmaniasis is a parasitic disease caused by the protozoan *Leishmania*, transmitted by female phlebotomine sandfly. The disease spectrum ranges from limited cutaneous plaques to disseminated nodules, and may involve viscera including liver, spleen and bone marrow. Concurrent occurrence of various patterns of the disease in a single patient is rare. Here, we report a patient with leishmania recidivans and mucocutaneous disease along with involvement of multiple bones, a combination that has not been reported in Pakistan.

Corresponding Author | Dr. Hira Tariq, Assistant Professor of Dermatology Department, SIMS/ Services Hospital, Lahore

Email: kemcolianhira46@gmail.com

Introduction:

Leishmaniasis is caused by the genus *Leishmania* which is an obligate, intramacrophage, protozoan parasite. The vector responsible for the disease is sandfly *Phlebotomus*. Along with the skin, the disease may involve mucous membranes and internal organs¹. Commonly effected parts of the world include South Asia and Middle East. Cutaneous leishmaniasis is further divided into classical cutaneous leishmaniasis, diffuse cutaneous leishmaniasis, leishmania recidivans and post Kala-azar dermal leishmaniasis². Visceral form of disease is of classical and viscerotropic type. Simultaneous occurrence of cutaneous and visceral forms of disease in a single host is seldom reported and is a rare entity³.


Case report

A 20 years old male from Hango, North of Pakistan, presented with recurrent multiple erythematous crusted plaques and nodules on face and limbs for last eight years. These plaques used to heal in a month to leave atrophic scars. Few plaques developed in old scars as

well. He developed multiple plaques on forehead, left cheek and left nasal mucosa, which also healed with scarring and deformity. He also developed nodule in middle finger of right hand with pain and restricted movement of interphalangeal joints. He had pain in multiple bones and joints especially those with overlying nodules, plaques and scars. He had weight loss and malaise. He had been on multiple medications including steroids, antimonials, antituberculous therapy and antibiotics, without relief. His sister too had similar lesions and was diagnosed as having ocular leishmaniasis.

On examination, he was well oriented, pale and was short for his age. He had hypertrichosis on face and body. He had reduced muscle mass on limbs. There were multiple crusted nodules and plaques on shoulders, knees and left foot (Figures 1-4). He had a firm subcutaneous nodule involving the middle finger of his right hand (Figure 3). It was immobile, tender and adherent to underlying structures. The movements of the finger were restricted. He had onychomycosis of multiple nails of his fingers and toes and pityriasis versicolor

on trunk and limbs. There were multiple atrophic depressed scars on face, shoulders, arms, hands, knees, legs and feet (Figure 5). Cervical lymph nodes were palpable. Many of his bones and joints were tender including shoulders, left knee and left shin. Rest of systemic examination was unremarkable and no visceromegaly was noted.



(Figure 1: Plaques on old scars on left shoulder.

Figure 2: Crusted plaque on right shoulder

Figure 3: Right hand showing onychomycosis of nails and subcutaneous nodule involving middle finger.

Figure 4: Crusted plaques on left knee


Figure 5: Old scars of healed lesions on face, involving underlying nasal mucosa)

His investigations showed hemoglobin of 9.4 g/dL, a total leucocyte count of 20,400/mm³, platelet count 713,000/mm³, an ESR of 70 with microcytic, hypochromic anemia, alkaline phosphatase was 764 U/L, LDH was 876 U/L, serum IgG level was 1775mg

/dL, anti-HIV, HBsAg and anti-HCV were negative. His sputum and smear from cutaneous lesions were also negative for tuberculous bacilli.

Smears taken from cutaneous nodules on shoulders demonstrated significant number of LD bodies when stained with Giemsa (Figure 6).

On histopathological examination of nodule in finger, there were numerous necrotizing granulomas. Most of them had lots of viable as well as degenerating neutrophils at the center. In addition, there were numerous LD bodies scattered all over. These were present within histiocytes as well as free within the tissue (Figure 7). Several degenerating bony spicules were also seen hinting at bone damage. Giemsa stain highlighted the LD bodies even further, which are seen as round to ovoid bodies measuring 3x1 to 6x3 μm (Figure 8).



Radiological investigations revealed soft tissue swellings and lytic lesions in bones along with obliteration of joint spaces, consistent with cutaneous and skeletal manifestations of leishmaniasis (Figures 9-



11). However, chest radiograph and abdominal ultrasound were normal.




Figure 9: Multiple lytic and degenerative lesions in the proximal and middle phalanges of the middle finger and carpal bones. Absence of periosteal reaction, marked osteopenia and reduced interdigital spaces along with soft tissue swelling around proximal interphalangeal joint of middle finger and increased soft tissue haze in lateral half of right hand. Multiple lytic lesions with marked osteopenia in bilateral distal femora (Figure 10) and skull (Figure 11)

Discussion

Based on the clinical features, other diagnostic considerations must include cutaneous tuberculosis, atypical mycobacterial infection, deep fungal infection, sarcoidosis and multiple myeloma. These were ruled out on histopathology, cultures and serology (Bence Jones proteins were also negative). Being an endemic area, cutaneous and visceral leishmaniasis has been reported from Pakistan.⁴ However, osseous involvement in the form of lytic lesions involving whole body has not been reported. Two patients have been reported to have osseous Leishmaniasis involving bones of distal extremities from Brazil⁵. Another patient with involvement of femur has been reported from Iran⁶. Our patient had involvement of multiple bones including skull and bones of limbs. Another interesting feature was presence of ocular leishmaniasis in the sibling. Leishmaniasis can have a myriad of histopathological features. The most common and diagnostic of these is granulomatous inflammation. These granulomas themselves can be of various types viz sarcoidal granulomas, necrotizing granulomas, suppurative granulomas and palisaded granulomas⁷. A mixture of patterns may be seen in a single case⁸. Our case showed necrotizing as well as suppurative granulomas along with bony spicu-

les. Leishmaniasis can affect and damage a variety of organs in the body⁹. The patient was immunocompromised as he had onychomycosis and pityriasis versicolor. Atypical forms of leishmaniasis have been reported in immunocompromised patient¹⁰. We demonstrated LD bodies along with bony spicules indicating bone involvement with leishmaniasis, a rarely reported entity. Another striking feature noted was presence of multiple patterns of disease in a single patient. This highlights the possibility of changing pattern of disease, microbial characteristics and ultimate manifestations in host.

Limitation

Facilities for Leishmanial culture are not available in Pakistan, therefore we couldn't know more about the species of the parasite involved.

Conclusion

Extensive work up in this case helped us to reach the diagnosis of cutaneous leishmaniasis with bone involvement. So, evaluation of bone lesions by radiographic examination seems to play an important role in identification and management of these lesions.

References

1. Mokni M. Leishmanioses cutanées [Cutaneous leishmaniasis]. *Ann Dermatol Venereol.* 2019;146(3):232-246.
2. Torres-Guerrero E, Quintanilla-Cedillo MR, Ruiz-Esmenjaud J, Arenas R. Leishmaniasis: a review. *F1000Res.* 2017;6(1):750.
3. Sundar S. Visceral leishmaniasis. *Trop Parasitol.* 2015;5(2):83-5.
4. Naveed SS, Raza N, BuxH, Firdous M, Rafi B. Aclinicohaematological study of visceral Leishmaniasis from northern Pakistan. *MC.* 2011;17(2):54-57.
5. Costa AA, Saldanha AC, Leite BM, Ramos B, Junior IA, Noronha AL, et al. Imaging exams of bone lesions in patients with diffuse cutaneous leishmaniasis (DCL). *Acta Trop.* 2005;96(1):9-15.
6. Sasannejad P, Mazloum Farsi Baf M, Zabolinejad N, Behdani R, Mazloum Farsi Baf M. Leishmaniasis with Femoral Bone Involvement. *J Coll Physicians Surg Pak.* 2016;26(6):

- 37-8.
7. Sundharkrishnan L, North JP. Histopathologic features of cutaneous leishmaniasis and use of CD1a staining for amastigotes in Old World and New World leishmaniasis. *J Cutan Pathol.* 2017;44(12):1005-1011.
8. Isaacs MJ, Hossler EW. Cutaneous leishmaniasis. *Cutis.* 2015;96(6):408-9.
9. Gawade S, Nanaware M, Gokhale R, Adhav P. Visceral leishmaniasis: A case report. *Australas Med J.* 2012;5(2):130-4.
10. Russo R, Laguna F, López-Vélez R, Medrano FJ, Rosenthal E, Cacopardo B, et al. Visceral leishmaniasis in those infected with HIV: clinical aspects and other opportunistic infections. *Ann Trop Med Parasitol.* 2003;97(1):99-105.