

## Research Article

# Incidence of COVID-19 Infection in Nurses and their Families: A Prospective Cohort Study in Lahore

Iram Manzoor,<sup>1</sup> Aqsa Hameed,<sup>2</sup> Momal Khan,<sup>3</sup> Allah Yar Khan,<sup>4</sup> Amna Ayub<sup>5</sup>

<sup>1-5</sup> Akhtar Saeed Medical and Dental College, Lahore.

### Abstract

**Background:** Nurses, as frontline workers, are at a higher risk of contracting COVID19 and serve as a potential source of transmitting this infection to their family members. Estimating risk in this population is critical for controlling the rate of secondary attack rate among their families.

**Objective:** To measure the incidence of COVID-19 in nurses and their families with associated risk factors for transmission of infection.

**Methods:** A prospective cohort study was conducted in Akhtar Saeed Medical and Dental college, Lahore and associated three hospitals. A cohort of 176 nurses was identified in March 2020 by using non-probability convenience sampling technique and was followed till September 2021. During this tenure of one and a half year, incidence was observed by following the cohort and using a self-structured questionnaire. With SPSS 23, Chi square test was applied and p value of < 0.05 was considered significant.

**Results:** In a cohort of 176 nurses, followed for one and a half year, incidence of COVID-19 was 37(21%). Secondary attack rate among household members of these nurses was 3%. A significant association was observed between the nursing staff and exposure to their family members ( $p = 0.002$ ). Vaccination rate in nurses at the start of the study was only 26%. Nurses who worked directly with COVID-19 patients reported high incidence ( $p = 0.009$ ). Nurses who used Personal protective equipment (PPE) were less affected ( $p = 0.003$ ). Nurses with comorbidities showed higher incidence ( $p = 0.02$ ). Rate of psychological distress was high ( $p = 0.007$ ).

**Conclusion:** Incidence of COVID-19 among nurses was 21% with secondary attack rate of 3% in families. Protective use of N 95 mask and comorbidities were associated with incidence.

**Corresponding Author** | Prof. Dr. Iram Manzoor, Director Medical Education/ Professor & HOD Community Medicine Department, Akhtar Saeed Medical and Dental College, Lahore. **Email:** iramdr123@yahoo.com

**Keywords** | Incidence, COVID-19, Nurses, Families, Secondary attack rate

### Introduction:

The outbreak of COVID-19 first emerged in Wuhan, Hubei Province, China, in December 2019 and the World Health Organization announced it as a public health emergency of global concern on January 30, 2020.<sup>1</sup> Coronavirus is spreading globally with 267,865,289 confirmed cases of COVID-19 and 5,285,888 deaths as per reported to WHO on 7th December, 2021.<sup>2</sup> Glo-

bally people of all nations, races, and socio-economic groups are affected but the health of healthcare workers is dramatically challenged.<sup>3</sup>

It has been estimated that in year 2020 almost 90,000 healthcare workers from all over world have been COVID positive and greater than 600 nurses have lost their lives.<sup>4</sup> According to regional analysis, a study in Wuhan showed that on March 26th 2020, total confirmed

cases among healthcare workers (HCW's) were 50006 out of which 52.06% were nurses while rest of them were doctors (33.62%) and other paramedical staff (14.33%).<sup>5</sup>

According to an Iranian study, COVID-19 has infected at least 90,000 healthcare workers until June 5<sup>th</sup>, 2020 and more than 260 nurses have died as a result of the pandemic.<sup>6</sup> As of June 29, 2021, Pakistan had reported a total of 0.95 million confirmed cases of COVID-19.<sup>7</sup> In Pakistan more than 440 cases have been reported in frontline healthcare workers out of which 8 confirmed cases reported death in different provinces by 2020. Out of these 440, 67(15.2%) were nurses, 216(49%) were doctors and 161(36.5%) were other healthcare workers<sup>4</sup> and the number is still increasing.

Several risk factors responsible for COVID-19 spread among healthcare workers are inadequate availability of personal protective equipment, work overload, poor infection prevention practices and pre-existing comorbidities, unavailability of diagnostic tests, exposure to infected patients and asymptomatic carriers.<sup>8</sup> Number of healthcare workers affected by COVID-19 is constantly increasing. This creates a lot of stress on nurses and their families.<sup>4</sup>

Healthcare providers should be carefully monitored from time to time because if they are infected, they can transmit it to their colleagues, healthy patients and family members.<sup>9</sup>

Nurses play a critical role in the treatment of COVID-19 patients and are more susceptible to infection. There is scarcity of literature on the incidence of COVID-19 among Pakistani Nurses. The objective of the study is to measure the frequency of COVID-19 infection in nurses and their families and associated risk factors for transmission of infection.

### Methods:

This is a prospective cohort study conducted with a cohort identification of 238 nurses, through non-probability, convenient sampling technique in Farooq Hospital West Wood Lahore, Akhtar Saeed Trust Teaching Hospital, Lahore and Farooq Hospital, Allama Iqbal Town Branch, Lahore. This cohort was identified on March 2020 and was followed till September 2021. Initially cohort consisted of 250 nurses but the final number of nurses remained in cohort was 176. Attrition

was observed due to non-willingness of participants to continue in research and change in vaccination status. Ethical approval of this research was taken from the Institutional review board of Akhtar Saeed Medical and Dental College, Lahore and Medical superintendents of the relevant hospitals before data collection from nurses. IRB letter number M-19/055/CM was originated to grant data collection to researchers. Before data collection, informed consent was taken from nurses in identified cohort. Data was collected by observations and interviews on self-structured questionnaire. This questionnaire was pretested through pilot study. The questionnaire comprised of three sections. The first section comprised of the sociodemographic information of nurses, which included age, sex, marital status, area of residence and family type of the participant nurses. The second section comprised of questions regarding Nurses' workplace, their access to PPE's contact with COVID19 patients, psychological distress faced during working in COVID wards, vaccination status, severity of COVID19 symptoms. In the third section, family members living in the same house as health care workers were asked about their vaccination status, contact history, onset of suspicious symptoms, comorbidities, treatment plan, hospitalization, and isolation status. Data was entered and analyzed in SPSS (Statistical Package for Social Sciences) version 23. Descriptive statistics like frequency and proportions were used for categorical variables. Mean and standard deviations were calculated for continuous variable. Data was presented in the form of frequency tables. Chi square test was applied and p value of less than 0.05 was considered significant. Secondary attack rate was calculated among relatives of nurses residing in the same place as these nurses. It was calculated as the total number of new cases emerging in household of nurses divided by the total number of households of these nurses. Primary cases were excluded from numerator and denominator both.

### Results:

The mean age of nurses in identified cohort was 26.4 ± 3.9 years. Majority of the participants 159(90%) were females. Those who belonged to nuclear family were 104 (59%). Incidence of COVID 19 was reported in 37(21%). Out of the total 176, 46(26%) got COVID vaccine, and maximum respondents 31(67.3%) had Sinopharm vaccine as shown in Table 1.

**Table 1: Sociodemographic Profile of Cohort of Nurses**

Age	Frequency (n)	Percentage (%)
20-30 years	153	86.9
31-40 years	21	11.9
41-50 years	2	1.1
<b>Gender</b>		
Male	17	9.7
Female	159	90.3
<b>Type of family</b>		
Nuclear	104	59.1
Extended	72	40.9
<b>Vaccination</b>		
Yes	46	26.1
No	130	73.9
<b>Type</b>		
Sinopharm	31	67.3
Sinovac	15	32.6
<b>Workplace</b>		
Akhtar Saeed Trust Teaching Hospital	44	25
Westwood Branch	84	47.7
Iqbal Town Branch	48	27.3
<b>Unit</b>		
ICU	31	17.6
COVID Isolation Ward	6	3.4
Emergency Unit	34	19.3
OPD	13	7.4
CCU	12	6.8
Any other Ward	80	45.5

**Table 2: Bivariate Analysis to Assess Risk Factors Associated with COVID-19 in Nurses**

Variables	Affected with COVID-19	Not effected	Total	P-Value
<b>Vaccination Status</b>				
Yes	10(21.7%)	36 (78.3%)	46	0.89
No	27(20.7%)	103 (79.2%)	130	
<b>Work with COVID-19 Patients</b>				
Yes	28(28%)	72 (72%)	100	
<b>Psychological Distress in COVID Ward</b>				
Yes	13(38.2%)	21 (61.8%)	34	0.00
No	15(22.7%)	51 (77.3%)	66	
<b>Use N-95</b>				
Yes	32(26.4%)	89 (73.6%)	121	0.02
No	0(0%)	5 (100%)	5	
<b>Comorbidities</b>				
Yes	8(40%)	12 (60%)	20	0.02
No	29(18.6%)	127 (81.4%)	156	
<b>Diabetes</b>				
Yes	4(40%)	6 (60%)	10	0.02
No	33(24.8%)	133 (75.2%)	166	

**Table 3: Bivariate Analysis to Assess Disease Pattern of COVID-19 in Nurses**

Variables	Affected with COVID-19	Not effected	Total	P-Value
<b>Test for COVID-19</b>				0.00
Yes	35(97.1%)	1 (2.8%)	36	
No	2(1.4%)	138(98.6%)	140	
<b>Source of Infection</b>				
Colleagues	2(100%)	0 (0%)	2	0.00
Patients	33(100%)	0 (0%)	33	
Friends	2(100%)	0 (0%)	2	
<b>Type of Test</b>				
<b>PCR</b>				
Antibody/Serology Test	13(100%)	0 (0%)	13	0.00
Chest Xray	7(100%)	0 (0%)	7	
HRCT	12(100%)	0 (0%)	12	
	4(100%)	0 (0%)	4	
<b>Symptoms you noted in case of being infected with COVID-19</b>				
Fever	17 (100%)	0 (0%)	17	0.00
Cough	6 (100%)	0 (0%)	6	
Sore Throat	7 (100%)	0 (0%)	7	
Fatigue	2 (100%)	0 (0%)	2	
Body Ache	1 (100%)	0 (0%)	1	
<b>Type of treatment in Symptomatic State</b>				
Self-Medication	14 (100%)	0 (0%)	14	0.00
Doctor Recommendation	17(100%)	0 (0%)	17	
Others	2(100%)	0 (0%)	2	
<b>Severity of COVID-19 Symptoms</b>				
Symptomless	4 (100%)	0 (0%)	4	0.00
Mild	20 (100%)	0 (0%)	20	
Moderate	7 (100%)	0 (0%)	7	
Severe	6 (100%)	0 (0%)	6	
<b>Most Affected System</b>				
GIT	4(100%)	0 (0%)	4	0.00
Respiratory	24(100%)	0 (0%)	24	
Circulatory System	9 (100%)	0 (0%)	9	
<b>Isolation</b>				
Yes	21(100%)	0 (0%)	21	0.00
NO	16(100%)	0 (0%)	16	
<b>Hospitalization</b>				
Yes	3(100%)	0 (0%)	3	0.00
No	34(100%)	0 (0%)	34	

Bivariate analysis showed that nurses got infection irrespective of their vaccination status ( $p=0.89$ ). Nurses who worked directly with COVID-19 patients had high incidence of disease ( $p=0.00$ ). Psychologically stress was reported while working in COVID ward ( $p=0.00$ ). Use of N-95 mask played a significant role in prevention of disease ( $p=0.02$ ). Comorbidities was seen in 20(11%) of the participants and majority 10(50%) reported diabetes ( $p=0.02$ ) as evident in Table 2.

Bivariate analysis showed significant association with positivity of PCR test and patients as a source of infection. Fever was the most significant symptom. Maximum number of patients had mild symptoms and took doctor's advice for treatment. Most affected system was respiratory system as shown in Table 3.

Severity associated with COVID-19 infection was noticed in 6(16%) of the respondents and 3(8%) of them got hospitalized. Mean drop in oxygen level along with standard deviation was  $87.3 \pm 11.01$ .

Mean number of family members residing at home was  $6.45 \pm 2.89$ . Total number of family members affected by COVID-19 were 35 and secondary attack rate was estimated by dividing the number of exposed persons developing disease within range of incubation period ( $n=35$ ) by total number of exposed persons in households ( $n=1136$ ) and multiplied by 100 and the calculated result was 3%. Mean number of family members with vaccination was  $0.81 \pm 1.480$  SD. Major comorbidity noticed in household members 7(58%) was hypertension.

### Discussion:

Results of this showed that incidence of COVID 19 was reported in 37 (21%) out of a cohort of 176 nurses. Another study conducted in Muscat, Oman reported that the majority of the health care workers affected with COVID-19 were nurses 77(38%).<sup>10</sup> Health care professionals, who have played a critical role in the response to COVID-19, maybe at a higher risk of contracting SARS COV-2 and then transferring it to their family, coworkers or both. According to current study the number of family members who got affected by COVID-19 were 35 with secondary attack rate of 3 % while another study showed (32.1%) of family members were IgG positive with Covid-19 infection.<sup>11</sup> A study in Italy showed that 33(9%) health care workers were believed to be the source of infection for family mem-

bers.<sup>12</sup>

Healthcare workers (HCW) are slated to be early recipients of SARS-CoV-2 vaccines due to increased risk of exposure to patients with COVID-19. Although nurses are front-line health workers, the vaccination status among them was inadequate 46(26.1%) only. Whereas a Chinese study found vaccination rate of nurses as 76.98%.<sup>13</sup> Results of this study showed that vaccination status was not associated with incidence of disease in nurses.

A significant association was found with incidence of COVID 19 in nurses with place of duty allocation. Another study conducted in Pakistan showed maximum number of cases in nurses working in emergency ward and isolation ward specific for COVID 19.<sup>14</sup>

Healthcare professionals and nurses experienced emotional stress in the fight against Covid-19 infection and 34(34%) of our respondents experienced it. A study conducted in Jordan revealed that 41% of nurses were psychologically affected by this pandemic.<sup>15</sup>

Results of this study showed use of personal protective equipment by 140(79.5%) of the nurses in identified cohort, another study conducted in Pakistan showed that 129(28%) of HCWS had adequate access to PPE.<sup>14</sup>

Majority of nurses 121 (96.03%) reported use of N 95 masks, which had significant for prevention of COVID 19 infection in this study. A study published in Critical care medicine in 2010 has already established the role in N95 masks in prevention of influenza and SAARS.<sup>16</sup> A recent study conducted in Pakistan showed that disposable face masks were used by 263(78%) of the health care providers during their hospital working hours.<sup>17</sup>

In this study high incidence of COVID 19 was reported in nurses with associated comorbidities, particularly diabetes. A survey conducted in an Indian hospital where similar comorbidities in health care workers associated with SARSCov-2 infection were reported as hypertension in (49%), diabetes mellitus (47%) and (5%) with ischemic heart disease.<sup>18</sup>

According to results of this study 36(20.5%) of the individuals had their COVID-19 test, 35.1% preferred PCR. Another study reveals that 32.3% of the nurses, were detected positive with Covid-19 infection through PCR in a hospital-based study.<sup>19</sup>

Majority 33 (89.1%) of the nurses in this cohort got

infected from their patients. Depending upon the severity of COVID-19 symptoms respondents in this study experienced mild 20(54%), moderate 7(18%) and severe 6(16%) symptoms. Similar results were reported in a study conducted in Egypt where severe illness was reported only in 8(11%) of the nurses.<sup>20</sup> Upon participants systemic inquiry it was found that 24(64.8%) respondents experienced respiratory tract symptoms. These results are in line with the study findings conducted in Wuhan.<sup>21</sup>

In this study 3(8%) of the respondents required hospitalization depending upon the severity while another study conducted among symptomatic health care workers in Washington, only 6(3%) reported hospitalization related to COVID-19.<sup>22</sup>

Being a prospective cohort study with a long follow-up of one and a half year is a biggest strength of this study in which researchers observed the true incidence of infection among nurses. but a longer duration of follow-up forced change of variables noted during observations particularly the vaccination status of nurses to counteract this factor, analysis of incidence of infection was based on initial data obtained at the beginning of cohort identification. The initial cohort was 238 with attrition rate of 6.2% the data was analyzed only for 176 retained nurses. Another major limitation was that the cohort of nurses was identified on the basis of convenient sampling based on 3 associated hospitals of Akhtar Saeed Medical and Dental College to make follow-up easier. Despite of all the limitations, this study reflects data from 3 different hospitals of Lahore so results can be generalized.

### Conclusion:

Incidence of COVID-19 among nurses was 21% with secondary attack rate of 3% in families. Statistically significant association was observed between incidence of COVID-19 infection among nurses and protective use of PPEs, comorbidities and psychological stress faced during COVID-19.

### Ethical Approval:

Given  
Conflict of Interest: The authors declare no conflict of interest.

### Funding Source:

None

### References:

- Lai X, Wang M, Qin C, Tan L, Ran L, Chen D et al. Coronavirus disease 2019 (COVID-2019) infection among health care workers and implications for prevention measures in a tertiary hospital in Wuhan, China. *JAMA network open*. 2020;3(5):e209666.
- World Health Organization. Weekly epidemiological update on COVID-19-7 December 2021. Available at: URL: <https://www.who.int/publications/m/item/weekly-epidemiological-update-on-covid-19---7-december-2021>.
- Sperling D. Ethical dilemmas, perceived risk, and motivation among nurses during the COVID-19 pandemic. *Nurs Ethics*. 2021;28(1):9–22.
- Meghani SM, Lalani LN. Critical care nurses as frontline warriors during covid-19 in Pakistan. *The World of Crit Care Nurs*. 2020;13(4):196–201.
- Zheng LZ, Wang XW, Zhou CZ, Liu QL, Li SL, Sun QS, et al. Analysis of the infection status of healthcare workers in wuhan during the covid-19 outbreak: A cross sectional study. *Clin. Infect. Dis*. 2020;71(16):2109–2113.
- Sabetian G, Moghadami M, Haghghi LH, Shahriarirad R, Fallahi MJ, Asmari N, et al. COVID-19 infection among healthcare workers: a cross-sectional study in southwest Iran. *Virology*. 2021;18(1):1–8.
- Qamar MA, Irfan O, Dhillon RA, Bhatti A, Sajid MI, Awan S, et al. Acceptance of COVID-19 vaccine in Pakistan: A nationwide cross-sectional study. *Cureus*. 2021;13(7):e16603.
- Ali SA, Noreen SN, Farooq IF, Bugshan AB, Vohra FV. Risk assessment of healthcare workers at frontline against covid 19. *Pak J Med Sci*. 2020;36(COVID19-S4):S99–103.
- Barranco RB, Ventura FV. Covid-19 and infection in healthcare workers: An emergency problem. *Med Leg J*. 2020;88(2):65–66.
- Al Maskari Z, Al Blushi A, Khamis F, Al Tai A, Al Salmi I, Al Harthi H, et al. Characteristics of healthcare workers infected with COVID-19: A cross-sectional observational study. *IJID*. 2021;102(6):32–6.
- Lorenzo D, Carrisi C. COVID-19 exposure risk for family members of healthcare workers: An observational study. *Int J Infect Dis*. 2020;98(4):287–9.
- Felice C, Di Tanna GL, Zanusi G, Grossi U. Impact of COVID-19 outbreak on healthcare workers in Italy: results from a national e-survey. *J. Community Health*. 2020;45(4):675–83.
- Wang MW, Wen W, Wang N, Zhou MY, Wang CY, Ni J, et al. COVID-19 Vaccination Acceptance Among Healthcare Workers and Nonhealthcare Workers in China: A Survey. 2021;9:709056.

14. Hakim M, Khattak FA, Muhammad S, Ismail M, Ullah N, Atiq Orakzai M, et al. Access and Use Experience of Personal Protective Equipment Among Frontline Healthcare Workers in Pakistan During the COVID-19 Emergency: A Cross-Sectional Study. *Health Secur.* 2021;19(2):140-9.
15. Shahrour G, Dardas LA. Acute stress disorder, coping self efficacy and subsequent psychological distress among nurses amid COVID19. *J. Nurs. Manag.* 2020; 28(7):1686-95.
16. Gralton J, McLaws ML. Protecting healthcare workers from pandemic influenza: N95 or surgical masks? *Crit Care Med.* 2010;38(2):657-67.
17. Dhahri AA, Iqbal MR, Khan AF. A cross-sectional survey on availability of facilities to healthcare workers in Pakistan during the COVID-19 pandemic. *Ann. Med. Surg.* 2020;59(2):127-30.
18. Mahajan NN, Mathe A, Patokar GA, Bahirat S, Lokhande PD, Rakh V, et al. Prevalence and clinical presentation of COVID-19 among healthcare workers at a dedicated hospital in India. *J Assoc Physicians India.* 2020;68(12):16-21.
19. Stock AD, Bader ER, Cezayirli P, Inocencio J, Chalmers SA, Yassari R, et al. COVID-19 infection among health-care workers: serological findings supporting routine testing. *Front. Med.* 2020;7(1):471.
20. El-Sokkary RH, El-Kholy A, Mohy Eldin S, Khater WS, Gad DM, Bahgat S, et al. Characteristics and predicting factors of Corona Virus Disease-2019 (COVID-19) among healthcare providers in a developing country. *Plos one.* 2021;16(1):e0245672.
21. Yifan T, Ying L, Chunhong G, Jing S, Rong W, Zhenyu L, et al. Symptom cluster of ICU nurses treating COVID-19 pneumonia patients in Wuhan, China. *J. Pain Symptom Manag.* 2020;60(1):48-53.
22. Mani NS, Budak JZ, Lan KF, Bryson-Cahn C, Zelikoff A, Barker GE, et al. Prevalence of coronavirus disease 2019 infection and outcomes among symptomatic healthcare workers in Seattle, Washington. *Clin. Infect. Dis.* 2020;71(10):2702-7.